

ACROSS-BREED EPD TABLES FOR THE YEAR 2006 ADJUSTED TO BREED DIFFERENCES FOR BIRTH YEAR OF 2004

L. D. Van Vleck and L. V. Cundiff

Roman L. Hruska U.S. Meat Animal Research Center, ARS, USDA, Lincoln and Clay Center, NE 68933

Introduction

This report is the year 2006 update of estimates of sire breed means from data of the Germplasm Evaluation (GPE) project at the U.S. Meat Animal Research Center (MARC) adjusted to a year 2004 base using EPD from the most recent national cattle evaluations. Factors to adjust EPD of 16 breeds to a common birth year of 2004 were calculated and are reported in Tables 1-3 for birth weight (BWT), weaning weight (WWT), and yearling weight (YWT) and in Table 4 for the MILK component of maternal weaning weight (MWWT).

There were few changes from the 2005 update (Van Vleck and Cundiff, 2005):

Records from USMARC for birth, weaning, and yearling weight were the same as last year. The EPD for the Tarentaise national cattle evaluations were updated this year which caused only minor changes in the across-breed adjustment factors for Tarentaise weights and maternal milk. Estimates of genetic parameters used for Shorthorn evaluations and resulting EPD have changed but across-breed adjustment factors for Shorthorn changed only slightly.

A considerable number of maternal records (weaning weights of grandprogeny) were added again this year, ranging from about 135 for Hereford and Angus to about 70 for Simmental, Limousin, Charolais, Gelbvieh, and Red Angus. Last year, for the first time, maternal records for Brangus (62) and Beefmaster (71) were included. Those numbers increased by about 60% to 99 and 113 this year.

- 1) For BWT, the MARC records were the same as last year so any slight changes from the analysis done last year will be due to EPD reported by the breed associations.
- 2) For WWT, the MARC records were also the same as last year. Most adjustment factors changed only slightly or not at all. The largest change was from 32.0 to 30.6 lb for Shorthorn.
- 3) For YWT, the same animals had records at MARC as last year. Adjustment factors relative to the Angus base for year 2004 changed little from those reported last year.
- 4) About 135 maternal weaning weights for both Hereford and Angus grandsires and about 70 for Simmental, Limousin, Charolais, Gelbvieh and Red Angus grandsires were added to the maternal analysis. Nearly 60% more maternal weaning weights were added for Brangus and Beefmaster grandsires. Changes in the across-breed

adjustments for MILK were not large but changed relatively more than for BWT, WWT, and YWT. The largest change (from -4.6 to -7.1) was for Beefmaster.

The across-breed table adjustments apply **only** to EPDs for most recent (in most cases; spring, 2006) national cattle evaluations. Serious biases can occur if the table adjustments are used with earlier EPDs which may have been calculated with a different within-breed base.

Materials and Methods

Adjustment for heterosis

The philosophy underlying the calculations has been that bulls compared using the across-breed adjustment factors will be used in a crossbreeding situation. Thus, calves and cows would generally exhibit 100% of both direct and maternal heterosis for the MILK analysis and 100% of direct heterosis for the BWT, WWT, and YWT analyses. The use of the MARC III composite (1/4 each of Pinzgauer, Red Poll, Hereford, and Angus) as a dam breed for Angus, Brangus, Hereford and Red Angus sires requires a small adjustment for level of heterozygosity for analyses of calves for BWT, WWT, and YWT and for cows for maternal weaning weight. Some sires (all multiple sire pasture mated) mated to the F1 cows are also crossbred so that adjustment for direct heterosis for the maternal analysis is required. Two approaches for accounting for differences in breed heterozygosity have been tried which resulted in similar final table adjustments. One approach was to include level of heterozygosity in the statistical models which essentially adjusts to a basis of no heterozygosity. The other approach was based on the original logic that bulls will be mated to another breed or line of dam so that progeny will exhibit 100% heterozygosity. Most of the lack of heterozygosity in the data results from homozygosity of Hereford or Angus genes from pure Hereford or Angus matings and also from Red Angus by Angus and from Hereford, Angus or Red Angus sires mated with MARC III composite dams. Consequently, the second approach was followed with estimates of heterosis obtained from analyses of BWT, WWT, YWT, and MWWT using only records from the imbedded diallel experiments with Hereford and Angus. Red Angus by Angus matings were assumed not to result in heterosis. With Brangus representing 5/8 and 3/8 inheritance from Angus and Brahman genes, records of Brangus sired calves were also adjusted to a full F1 basis when dams were Angus cows and MARC III cows (1/4 Angus). The adjustment for calves with Beefmaster (1/2 Brahman, 1/4 Shorthorn, 1/4 Hereford) sires was only when dams were MARC III cows (1/4 Hereford) as Beefmaster sires were not mated to Hereford cows.

The steps were:

- 1) Analyze records from Hereford-Angus (H-A) diallel experiments to estimate direct heterosis effects for BWT, WWT, YWT (1,326, 1,279, and 1,249 records for BWT, WWT, and YWT, respectively, representing 152 sires). The H-A diallel experiments were conducted as part of Cycle I (1970-1972 calf crops), Cycle II (1973-1974), Cycle IV (1986-1990) and Cycle VII (1999-2001) of the GPE program at MARC.

- 2) Adjust maternal weaning weight (MWWT) records of calves of the Hereford and Angus cows from the diallel for estimates of direct heterosis from Step 1) and then estimate maternal heterosis effects from 3,404 weaning weight records of 795 daughters representing 174 Hereford and Angus maternal grandsires.
- 3) Adjust all records used for analyses of BWT, WWT, and YWT for lack of direct heterozygosity using estimates from Step 1), and
- 4) Adjust all records used for analysis of MWWT for lack of both direct and maternal heterozygosity using estimates from Steps 1) and 2).

Models for the analyses to estimate heterosis were the same as for the across-breed analyses with the obvious changes in breed of sire and breed of dam effects. Estimates of direct heterosis were 3.01, 14.70, and 30.39 lb for BWT, WWT, and YWT, respectively. The estimate of maternal heterosis was 23.81 lb for MWWT. As an example of step 3), birth weight of a Hereford by Hereford calf would have 3.01 added. A Red Angus by MARC III calf would have $(1/4)$ (3.01) added to its birth weight. A Red Poll sired calf of an Angus by MARC III dam would have $(1/8)$ (14.70) plus $(1/4)$ (23.20) added to its weaning weight record to adjust to 100% heterozygosity for both direct and maternal components of weaning weight.

After these adjustments, all calculations were as outlined in the 1996 BIF Guidelines. The basic steps were given by Notter and Cundiff (1991) with refinements by Núñez-Dominguez et al. (1993), Cundiff (1993, 1994), Barkhouse et al. (1994, 1995), and Van Vleck and Cundiff (1997–2005). All calculations were done with programs written in Fortran language with estimates of variance components, regression coefficients, and breed effects obtained using the MTDFREML package (Boldman et al., 1995). All breed solutions are reported as differences from Angus. The table values of adjustment factors to add to within-breed EPD are relative to Angus.

Models for Analysis of MARC Records

Fixed effects in the models for BWT, WWT (205-d), and YWT (365-d) were: breed of sire (17 including Pinzgauer); dam line (Hereford, Angus, selection lines of Herefords, MARC III composite) by sex (female, male) by age of dam (2, 3, 4, 5-9, ≥ 10 yr) combination (49), year of birth (21) of dam (1970-76, 86-90, 92-94, 97-99, and 2000-02) by damline combination (101) and a separate covariate for day of year at birth of calf for each of the three breeds of dam. Cows from the Hereford selection lines were used in Cycle IV of GPE. To account for differences from the original Hereford cows, Hereford dams were subdivided into the selection lines and others. That refinement of the model had little effect on breed of sire solutions. Dam of calf was included as a random effect to account for correlated maternal effects for cows with more than one calf (4,630 dams for BWT, 4,395 for WWT, 4,243 for YWT). For estimation of variance components and to estimate breed of sire effects, sire of calf was also used as a random effect (650).

Variance components were estimated with a derivative-free REML algorithm. At convergence, the breed of sire solutions were obtained as were the sampling variances

of the estimates to use in constructing prediction error variances for pairs of bulls of different breeds.

For estimation of coefficients of regression of progeny performance on EPD of sire the random sire effect was dropped from the model. Pooled regression coefficients, and regression coefficients by sire breed, by dam line, and by sex of calf were obtained. These regression coefficients are monitored as accuracy checks and for possible genetic by environment interactions. The pooled regression coefficients were used as described in the next section to adjust for genetic trend and bulls used at MARC.

The fixed effects for the analysis of maternal effects included breed of maternal grandsire (17 including Pinzgauer), maternal grandam line (Hereford, Angus, MARC III), breed of natural service mating sire (17), sex of calf (2), birth year-GPE cycle-age of dam subclass (87), and mating sire breed by GPE cycle by age of dam subclass (46) with a covariate for day of year of birth. The subclasses are used to account for confounding of years, mating sire breeds, and ages of dams. Age of dam classes were 2, 3, 4, 5-9, ≥ 10 yr. For estimation of variance components and estimation of breed of maternal grandsire effects, random effects were maternal grandsire (619) and dam (3,214 daughters of the maternal grandsires). Mating sires were unknown within breed. For estimation of regression coefficients of grandprogeny weaning weight on maternal grandsire EPD for weaning weight and milk, random effects of both maternal grandsire and dam (daughter of MGS) were dropped from the model.

Adjustment of MARC Solutions

The calculations of across-breed adjustment factors rely on solutions for breed of sire or breed of maternal grandsire from records at MARC and on averages of within-breed EPD from the breed associations. The records from MARC are not used in calculation of within-breed EPD by the breed associations. The basic calculations for BWT, WWT, and YWT are as follows:

MARC breed of sire solution for breed i (MARC (i)) adjusted for genetic trend (as if bulls born in the base year had been used rather than the bulls actually used):

$$M_i = \text{MARC } (i) + b[\text{EPD}(i)_{YY} - \text{EPD}(i)_{\text{MARC}}].$$

Breed Table Factor (A_i) to add to the EPD for a bull of breed i :

$$A_i = (M_i - M_x) - (\text{EPD}(i)_{YY} - \text{EPD}(x)_{YY})$$

where,

MARC(i) is solution for sire breed i from mixed model equations with MARC data,

EPD(i)_{YY} is the average within-breed EPD for breed i for animals born in the base year (YY, which is two years before the update; e.g., YY = 2004 for the 2006 update),

EPD(i)_{MARC} is the weighted (by number of progeny at MARC) average of EPD of bulls of breed i having progeny with records at MARC,

b is the pooled coefficient of regression of progeny performance at MARC on EPD of sire (for 2006: 1.02, 0.89, and 1.14 for BWT, WWT, YWT),

i denotes sire breed i, and

x denotes the base breed, which is Angus in this report.

The calculations to arrive at the Breed Table Factor for MILK are more complicated because of the need to separate the direct effect of the maternal grandsire breed from the maternal effect of the breed.

MARC breed of maternal grandsire solution for WWT adjusted for genetic trend:

$$\begin{aligned} \text{MWWT}(i) = & \text{MARC}(i)_{\text{MGS}} + b_{\text{wwt}}[\text{EPD}(i)_{\text{YYWWT}} - \text{EPD}(i)_{\text{MARCWWT}}] \\ & + b_{\text{MLK}}[\text{EPD}(i)_{\text{YYMLK}} - \text{EPD}(i)_{\text{MARCMLK}}] \end{aligned}$$

MARC breed of maternal grandsire solution (MWWT(i)) adjusted for genetic trend and direct genetic effect:

$$\text{MILK}(i) = [\text{MWWT}(i) - 0.5 \text{M}(i)] - [\text{ } - 0.5 \bar{\text{M}}]$$

Breed Table Factor to add to EPD for MILK for bull of breed i:

$$A_i = [\text{MILK}(i) - \text{MILK}(x)] - [\text{EPD}(i)_{\text{YYMLK}} - \text{EPD}(i)_{\text{MARCMLK}}]$$

where,

MARC(i)_{MGS} is solution for MGS breed i for WWT from mixed model equations with MARC data,

EPD(i)_{YYWWT} is the average within-breed EPD for WWT for breed i for animals born in base year (YY),

EPD(i)_{MARCWWT} is the weighted (by number of grandprogeny at MARC) average of EPD for WWT of MGS of breed i having grandprogeny with records at MARC,

EPD(i)_{YYMLK} is the average within-breed EPD for MILK for breed i for animals born in base year (YY),

EPD(i)_{MARCMLK} is the weighted (by number of grandprogeny at MARC) average of EPD for MILK of MGS of breed i having grandprogeny with records at MARC,

b_{WWT} , b_{MLK} are the coefficients of regression of performance of MARC grandprogeny on MGS EPD for WWT and MILK (for 2006: 0.57 and 1.13),

$M(i) = M_i$ is the MARC breed of sire solution from the first analysis of direct breed of sire effects for WWT adjusted for genetic trend,

\overline{MWWT} and \overline{M} are unneeded constants corresponding to unweighted averages of $MWWT(i)$ and $M(i)$ for $i = 1, \dots, n$, the number of sire (maternal grandsire) breeds included in the analysis.

Results

Tables 1, 2, and 3 (for BWT, WWT, and YWT) summarize the data from, and results of, MARC analyses to estimate breed of sire differences and the adjustments to the breed of sire effects to a year 2004 base. The last column of each table corresponds to the Breed Table Factor for that trait.

The general result shown in Tables 1-4 is that for weaning weight many breeds are continuing to become more similar to the arbitrary base breed, Angus. For yearling weight, however, Angus is becoming heavier in comparison to differences in the 2005 report. Most of the other breeds have not changed much relative to each other. Column 7 of Tables 1-3 and column 10 of Table 4 represent the best estimates of breed differences for calves born in 2004. These pairs of differences minus the corresponding differences in average EPD for animals born in 2004 result in the last column of the tables to be used as adjustment factors for pairs of sires having within-breed EPD.

Birth Weight

The range in estimated breed of sire differences for BWT relative to Angus is large: from 1.1 lb for Red Angus to 9.0 lb for Charolais and 12.0 lb for Brahman. The relatively heavy birth weights of Brahman sired progeny would be expected to be completely offset by favorable maternal effects reducing birth weight if progeny were from Brahman or Brahman cross dams which would be an important consideration in crossbreeding programs involving Brahman cross females. Differences from Angus were only slightly changed from the 2005 update (Van Vleck and Cundiff, 2005).

Suppose the EPD for birth weight for a Charolais bull is +2.0 (which is above the year 2004 average of 1.3 for Charolais) and for a Hereford bull is also +2.0 (which is below the year 2004 average of 3.7 for Herefords). The across-breed adjustment factors in the last column of Table 1 are 2.9 for Hereford and 10.0 for Charolais. Then the adjusted EPD for the Charolais bull is $10.0 + 2.0 = 12.0$ and for the Hereford bull is $2.9 + 2.0 = 4.9$. The expected birth weight difference when both are mated to another breed of cow, e.g., Angus, would be $12.0 - 4.9 = 7.1$ lb.

Weaning Weight

Weaning weights have continued to become more similar to Angus for most (10) breeds but slightly smaller for five breeds than for Angus for use as sire breeds. Most of

the changes between the year 2005 and 2006 updates were less than 2 lb. All except four sire breed means for WWT adjusted to year of birth of 2004 are within about 10 lb of the Angus mean.

Yearling Weight

Adjusted to a base year of 2004, Angus had heavier yearling weights than 10 breeds (12.6 to 53.6 lb), slightly heavier weights than 3 breeds (3.6 to 5.9 lb), and lighter yearling weights than 2 breeds (8.8 and 16.8 lb).

MILK

The changes from last year for MILK compared to Angus for the current base year were generally small.

Accuracies and Variance Components

Table 5 summarizes the average Beef Improvement Federation (BIF) accuracy for bulls with progeny at MARC weighted appropriately by number of progeny or grandprogeny. South Devon bulls had relatively small accuracy for all traits as did Hereford, Brahman, and Maine-Anjou bulls. Braunvieh bulls had low accuracy for milk. The accuracy values for Brangus are relatively high. Table 6 reports the estimates of variance components from the records that were used in the mixed model equations to obtain breed of sire and breed of MGS solutions. Neither Table 5 nor Table 6 changed much from the 2004 report.

Regression Coefficients

Table 7 updates the coefficients of regression of records of MARC progeny on sire EPDs for BWT, WWT, and YWT which have theoretical expected values of 1.00. The standard errors of the specific breed regression coefficients are large relative to the regression coefficients. Large differences from the theoretical regressions, however, may indicate problems with genetic evaluations, identification, or sampling. The pooled (overall) regression coefficients of 1.02 for BWT, 0.89 for WWT, and 1.14 for YWT were used to adjust breed of sire solutions to the base year of 2004. These regression coefficients are reasonably close to expected values of 1.0. Deviations from 1.00 are believed to be due to scaling differences between performance of progeny in the MARC herd and of progeny in herds contributing to the national genetic evaluations of the 16 breeds.

The regression coefficient for female progeny on sire EPDs for YWT was 1.00 compared to 1.28 for steers. These differences might be expected because postweaning average daily gains for heifers at MARC have been significantly less than those for steers. The heifers were fed relatively high roughage diets to support average daily gains of 1.6 lb per day while the steers were fed relatively high energy growing and finishing diets supporting average daily gains of about 3.4 lb per day. For reasons that

have never been clear, the regressions for sex used to fluctuate widely from year to year, but for the past few years the pattern has been fairly consistent (female estimates have ranged from 0.93 to 1.02; while male estimates have ranged from 1.26 to 1.32).

The coefficients of regression of records of grandprogeny on MGS EPDs for WWT and MILK are shown in Table 8. Several sire (MGS) breeds have regression coefficients considerably different from the theoretical expected values of 0.50 for WWT and 1.00 for MILK. Standard errors, however, for the regression coefficients by breed are large except for Angus and Hereford. The standard errors for regression coefficients over all breeds of grandsires associated with heifers and steers overlap for MILK EPDs. The pooled regression coefficients of 0.57 for MWWT and 1.13 for MILK are reasonably close to the expected regression coefficients of 0.50 and 1.00.

Prediction Error Variances of Across-Breed EPD

The standard errors of differences in the solutions for breed of sire and breed of MGS differences from the MARC records can be adjusted by theoretical approximations to obtain variances of adjusted breed differences (Van Vleck, 1994; Van Vleck and Cundiff, 1994). These variances of estimated breed differences can be added to prediction error variances of within-breed EPDs to obtain prediction error variances (PEV) or equivalently standard errors of prediction (SEP) for across-breed EPDs (Van Vleck and Cundiff, 1994, 1995). The variances of adjusted breed differences are given in the upper triangular part of Table 9 for BWT, lower triangular part of Table 9 for YWT, upper triangular part of Table 10 for direct WWT, and lower triangular part of Table 10 for MILK. Use of these tables to calculate standard errors of prediction for expected progeny differences of pairs of bulls of the same or different breeds was discussed in the 1995 BIF proceedings (Van Vleck and Cundiff, 1995).

Even though the variances of estimates of adjusted breed differences look large, especially for YWT and MILK, they generally contribute a relatively small amount to standard errors of predicted differences. For example, suppose for WWT, a Salers bull has an EPD of 15.0 with prediction error variance of 75 (SEP = 8.7) and a Hereford bull has an EPD of 30.0 with PEV of 50 (SEP = 7.1). The difference in predicted progeny performance is (Salers adjustment + Salers bull's EPD) - (Hereford adjustment + Hereford bull's EPD):

$$(29.0 + 15.0) - (-2.5 + 30.0) = 44.0 - 27.5 = 16.5.$$

The prediction error variance for this difference is (use the 18.0 in the upper part of Table 10 at intersection of row for HE and column for SA):

$V(\text{Salers breed} - \text{Hereford breed}) + \text{PEV}(\text{Salers bull}) + \text{PEV}(\text{Hereford bull})$:

$$18 + 75 + 50 = 143$$

with

standard error of prediction, $SEP = \text{SQRT}(143) = 12$.

If the difference between the Salers and Hereford breeds in the year 2004 could be estimated perfectly, the variance of the estimate of the breed difference would be 0 and the standard error of prediction between the two bulls would be:

$SEP(\text{difference}) = \text{SQRT}(0 + 75 + 50) = 11.2$ which is only slightly smaller than 12.0.

Implications

Bulls of different breeds can be compared on a common EPD scale by adding the appropriate table factor to EPDs produced in the most recent genetic evaluations for each of the 16 breeds. The across-breed EPDs are most useful to commercial producers purchasing bulls of two or more breeds to use in systematic crossbreeding programs. Uniformity in across-breed EPDs should be emphasized for rotational crossing. Divergence in across-breed EPDs for direct weaning weight and yearling weight should be emphasized in selection of bulls for terminal crossing. Divergence favoring lighter birth weight may be helpful in selection of bulls for use on first calf heifers. Accuracy of across-breed EPDs depends primarily upon the accuracy of the within-breed EPDs of individual bulls being compared.

References

- Barkhouse, K. L., L. D. Van Vleck, and L. V. Cundiff. 1994. Breed comparisons for growth and maternal traits adjusted to a 1992 base. Proc. BIF Research Symposium, Des Moines, IA, May, 1994. pp 197-209.
- Barkhouse, K. L., L. D. Van Vleck, and L. V. Cundiff. 1995. Mixed model methods to estimate breed comparisons for growth and maternal traits adjusted to a 1993 base. Proc. BIF Research Symposium, Sheridan, WY. May 31-June 3, 1995. pp 218-239.
- Boldman, K. G., L. A. Kriese, L. D. Van Vleck, and S. D. Kachman. 1993. *A Manual for Use of MTDFREML* (DRAFT). A set of programs to obtain estimates of variances and covariances. USDA-ARS, Roman L. Hruska U.S. Meat Animal Research Center, Clay Center, NE. (120 pp).
- Cundiff, L. V. 1993. Breed comparisons adjusted to a 1991 basis using current EPD's. Proc. BIF Research Symposium, Asheville, NC, May 26-29, 1993. pp 114-123.
- Cundiff, L. V. 1994. Procedures for across breed EPD's. Proc. Fourth Genetic Prediction Workshop, Beef Improvement Federation, Kansas City, MO, Jan. 1994.
- Notter, D. R., and L. V. Cundiff. 1991. Across-breed expected progeny differences: Use of within-breed expected progeny differences to adjust breed evaluations for sire sampling and genetic trend. J. Anim. Sci. 69:4763-4776.
- Núñez-Dominguez, R., L. D. Van Vleck, and L. V. Cundiff. 1993. Breed comparisons for growth traits adjusted for within-breed genetic trend using expected progeny differences. J. Anim. Sci. 71:1419-1428.
- Van Vleck, L. D., and L. V. Cundiff. 1994. Prediction error variances for inter-breed genetic evaluations. J. Anim. Sci. 71:1971-1977.
- Van Vleck, L. D., and L. V. Cundiff. 1995. Assignment of risk to across-breed EPDs with tables of variances of estimates of breed differences. Proc. BIF Research Symposium, Sheridan, WY. May 31-June 3, 1995. pp 240-245.
- Van Vleck, L. D., and L. V. Cundiff. 1997. The across-breed EPD tables adjusted to a 1995 base. Proc. BIF Research Symposium, Dickinson, ND. May 14-17, 1997. pp 102-117.
- Van Vleck, L. D., and L. V. Cundiff. 1997. Differences in breed of sire differences for weights of male and female calves. Proc. BIF Research Symposium, Dickinson, ND. May 14-17, 1997. pp 131-137.

Van Vleck, L. D., and L. V. Cundiff. 1998. Across-breed EPD tables for 1998 adjusted to a 1996 base. Proc. Beef Improvement Federation Research Symposium and Annual Meeting. Calgary, Alberta, Canada. July 2, 1998. pp 196-212.

Van Vleck, L. D. 1994. Prediction error variances for inter-breed EPD's. Proc. Fourth Genetic Prediction Workshop, Beef Improvement Federation, Kansas City, MO, Jan. 1994.

Van Vleck, L. D., and L. V. Cundiff. 1998. Influence of breed of dam on across-breed adjustment factors. Midwestern Section ASAS and Midwest Branch ADSA 1998 Meeting, Des Moines, IA. Abstract # 10. p 31.

Van Vleck, L. D., and L. V. Cundiff. 1999. Across-breed EPD tables for 1999 adjusted to a 1997 base. Proc. Beef Improvement Federation Research Symposium and Annual Meeting, Roanoke, VA. June 15-19, 1999. pp 155-171.

Van Vleck, L. D., and L. V. Cundiff. 2000. Across-breed EPD tables for 2000 adjusted to a 1998 base. Proc. Beef Improvement Federation Research Symposium and Annual Meeting, Wichita, KS. July 12-15, 2000. pp 98-116.

Van Vleck, L. D., and L. V. Cundiff. 2001. Across-breed EPD tables for 2001 adjusted to breed differences for birth year 1999. Proc. Beef Improvement Federation Research Symposium and Annual Meeting, San Antonio, TX. July 11-14, 2001. pp 44-63

Van Vleck, L. D., and L. V. Cundiff. 2002. Across-breed EPD tables for 2002 adjusted to breed differences for birth year of 2000. Proc. Beef Improvement Federation Research and Annual Meeting, Omaha, NE. July 10-13, 2002. pp 139-159.

Van Vleck, L. D., and L. V. Cundiff. 2003. Across-breed EPD tables for the year 2003 adjusted to breed differences for birth year of 2001. Beef Improvement Federation Research and Annual Meeting, Lexington, KY. May 28-31, 2003. pp 55-63.

Van Vleck, L. D., and L. V. Cundiff. 2004. Across-breed EPD tables for the year 2004 adjusted to breed differences for birth year of 2002. Beef Improvement Federation Research and Annual Meeting, Sioux Falls, SD. May 25-28, 2004. pp 46-61.

Van Vleck, L. D., and L. V. Cundiff. 2005. Across-breed EPD tables for the year 2005 adjusted to breed differences for birth year of 2003. Beef Improvement Federation Research and Annual Meeting, Billings, MT. July 6-9, 2005. pp 126-142.

Table 1. Breed of sire solutions from MARC, mean breed and MARC EPD used to adjust for genetic trend to the year 2004 base and factors to adjust within breed EPD to an Angus equivalent – BIRTH WEIGHT (lb)

| Breed | Number | | Raw | Ave. Base EPD | | Breed Soln | | Adjust to | | Factor to |
|-------------|--------|---------|---------------|----------------|----------------|------------------|-------------|------------------|---------------|-------------------------|
| | Sires | Progeny | MARC Mean (1) | Breed 2004 (2) | MARC Bulls (3) | + Ang vs Ang (4) | at MARC (5) | + Ang vs Ang (6) | 2004 Base (7) | adjust EPD To Angus (8) |
| Hereford | 113 | 1817 | 87 | 3.7 | 2.4 | 88 | 3.6 | 89 | 4.3 | 2.9 |
| Angus | 105 | 1421 | 84 | 2.3 | 1.7 | 84 | 0.0 | 85 | 0.0 | 0.0 |
| Shorthorn | 25 | 181 | 87 | 1.8 | 1.0 | 90 | 6.4 | 91 | 6.6 | 7.1 |
| South Devon | 15 | 153 | 80 | 0.2 | -0.1 | 88 | 4.3 | 89 | 4.0 | 6.1 |
| Brahman | 40 | 589 | 98 | 1.8 | 0.7 | 96 | 11.6 | 97 | 12.0 | 12.5 |
| Simmental | 48 | 623 | 87 | 1.8 | 2.8 | 91 | 7.0 | 90 | 5.3 | 5.8 |
| Limousin | 40 | 589 | 83 | 2.1 | 0.6 | 87 | 3.0 | 89 | 3.9 | 4.1 |
| Charolais | 75 | 675 | 89 | 1.3 | 0.4 | 93 | 8.8 | 94 | 9.0 | 10.0 |
| Maine-Anjou | 18 | 218 | 94 | 2.5 | 5.9 | 95 | 10.6 | 91 | 6.5 | 6.3 |
| Gelbvieh | 48 | 595 | 89 | 1.9 | 1.2 | 88 | 4.1 | 89 | 4.3 | 4.7 |
| Tarentaise | 7 | 199 | 80 | -1.5 | 1.7 | 87 | 3.2 | 87 | 2.3 | 3.1 |
| Salers | 27 | 189 | 85 | 1.1 | 1.8 | 88 | 4.4 | 88 | 3.0 | 4.2 |
| Red Angus | 21 | 206 | 85 | 0.4 | -0.8 | 85 | 0.6 | 86 | 1.1 | 3.0 |
| Braunvieh | 7 | 188 | 88 | 1.1 | 0.8 | 89 | 5.1 | 89 | 4.8 | 6.0 |
| Brangus | 21 | 215 | 91 | 2.0 | 2.4 | 90 | 5.9 | 89 | 4.9 | 5.2 |
| Beefmaster | 21 | 214 | 96 | 0.4 | 0.8 | 92 | 8.3 | 92 | 7.3 | 9.2 |

Calculations:

$$(4) = (5) + (1, \text{Angus})$$

$$(6) = (4) + b[(2) - (3)] \text{ with } b = 1.02$$

$$(7) = (6) - (6, \text{Angus})$$

$$(8) = (7) - (7, \text{Angus}) - [(2) - (2, \text{Angus})]$$

Table 2. Breed of sire solutions from MARC, mean breed and MARC EPD used to adjust for genetic trend to the year 2004 base and factors to adjust within breed EPD to an Angus equivalent – WEANING WEIGHT (lb)

| Breed | Number | | Raw | Ave. Base EPD | | Breed Soln | | Adjust to | | Factor to |
|-------------|--------|---------|---------------|----------------|----------------|------------------|-------------|------------------|---------------|-------------------------|
| | Sires | Progeny | MARC Mean (1) | Breed 2004 (2) | MARC Bulls (3) | + Ang vs Ang (4) | at MARC (5) | + Ang vs Ang (6) | 2004 Base (7) | adjust EPD To Angus (8) |
| Hereford | 112 | 1712 | 503 | 37.0 | 23.5 | 501 | -2.7 | 513 | -4.0 | -2.5 |
| Angus | 106 | 1315 | 504 | 38.5 | 23.5 | 504 | 0.0 | 517 | 0.0 | 0.0 |
| Shorthorn | 25 | 170 | 521 | 13.0 | 8.1 | 518 | 14.1 | 522 | 5.1 | 30.6 |
| South Devon | 15 | 134 | 443 | 19.1 | -0.1 | 503 | -0.6 | 521 | 3.2 | 22.6 |
| Brahman | 40 | 509 | 532 | 14.1 | 4.3 | 520 | 16.1 | 529 | 11.5 | 35.9 |
| Simmental | 47 | 564 | 505 | 34.1 | 23.8 | 526 | 22.4 | 536 | 18.2 | 22.6 |
| Limousin | 40 | 533 | 477 | 36.3 | 20.8 | 503 | -0.8 | 517 | -0.4 | 1.8 |
| Charolais | 74 | 600 | 514 | 20.0 | 8.4 | 527 | 23.3 | 538 | 20.3 | 38.8 |
| Maine-Anjou | 18 | 197 | 459 | 39.6 | 47.3 | 519 | 15.1 | 512 | -5.1 | -6.2 |
| Gelbvieh | 48 | 559 | 507 | 41.0 | 32.3 | 518 | 14.3 | 526 | 8.7 | 6.2 |
| Tarentaise | 7 | 191 | 476 | 4.0 | -4.7 | 507 | 2.7 | 515 | -2.8 | 31.7 |
| Salers | 27 | 176 | 525 | 15.5 | 7.0 | 516 | 11.7 | 523 | 6.0 | 29.0 |
| Red Angus | 21 | 199 | 535 | 29.0 | 27.6 | 505 | 1.0 | 506 | -11.1 | -1.6 |
| Braunvieh | 7 | 183 | 451 | 7.0 | 7.3 | 516 | 12.0 | 516 | -1.6 | 29.9 |
| Brangus | 21 | 208 | 550 | 22.7 | 25.9 | 524 | 20.3 | 521 | 4.1 | 19.9 |
| Beefmaster | 22 | 215 | 563 | 7.0 | 13.7 | 530 | 26.3 | 524 | 7.0 | 38.5 |

Calculations:

$$(4) = (5) + (1, \text{Angus})$$

$$(6) = (4) + b[(2) - (3)] \text{ with } b = 0.89$$

$$(7) = (6) - (6, \text{Angus})$$

$$(8) = (7) - (7, \text{Angus}) - [(2) - (2, \text{Angus})]$$

Table 3. Breed of sire solutions from MARC, mean breed and MARC EPD used to adjust for genetic trend to the year 2004 base and factors to adjust within breed EPD to an Angus equivalent – YEARLING WEIGHT (lb)

| Breed | Number | | Raw | Ave. Base EPD | | Breed Soln | | Adjust to | | Factor to |
|-------------|--------|---------|---------------|----------------|----------------|------------------|--------------------------|----------------------------|-------|-------------------------|
| | Sires | Progeny | MARC Mean (1) | Breed 2004 (2) | MARC Bulls (3) | + Ang vs Ang (4) | at MARC + Ang vs Ang (5) | 2004 Base + Ang vs Ang (6) | (7) | adjust EPD To Angus (8) |
| Hereford | 112 | 1627 | 852 | 63.0 | 39.7 | 852 | -20.0 | 879 | -24.2 | -15.7 |
| Angus | 106 | 1257 | 872 | 71.5 | 44.6 | 872 | 0.0 | 903 | 0.0 | 0.0 |
| Shorthorn | 25 | 168 | 918 | 21.0 | 12.3 | 887 | 15.0 | 897 | -5.9 | 44.6 |
| South Devon | 15 | 134 | 744 | 26.4 | -0.4 | 868 | -3.7 | 899 | -3.8 | 41.3 |
| Brahman | 40 | 438 | 838 | 23.1 | 7.9 | 832 | -40.1 | 849 | -53.6 | -5.2 |
| Simmental | 47 | 528 | 852 | 59.5 | 39.5 | 889 | 16.7 | 912 | 8.8 | 20.8 |
| Limousin | 40 | 527 | 797 | 68.2 | 42.5 | 849 | -23.3 | 878 | -24.8 | -21.5 |
| Charolais | 74 | 566 | 882 | 35.2 | 15.5 | 897 | 25.1 | 920 | 16.8 | 53.1 |
| Maine-Anjou | 18 | 196 | 787 | 78.2 | 94.1 | 884 | 12.3 | 866 | -36.7 | -43.4 |
| Gelbvieh | 48 | 555 | 849 | 73.0 | 57.7 | 864 | -7.8 | 882 | -21.1 | -22.6 |
| Tarentaise | 7 | 189 | 807 | 11.0 | -3.9 | 837 | -35.2 | 854 | -49.0 | 11.5 |
| Salers | 27 | 173 | 898 | 25.8 | 8.8 | 880 | 7.8 | 899 | -3.6 | 42.1 |
| Red Angus | 21 | 194 | 916 | 51.0 | 47.4 | 877 | 5.4 | 882 | -21.3 | -0.8 |
| Braunvieh | 7 | 182 | 737 | 8.0 | 11.8 | 856 | -16.4 | 851 | -51.6 | 11.9 |
| Brangus | 21 | 152 | 977 | 37.8 | 43.0 | 896 | 24.1 | 890 | -12.6 | 21.1 |
| Beefmaster | 22 | 157 | 991 | 12.0 | 22.7 | 893 | 20.9 | 881 | -22.2 | 37.3 |

Calculations:

$$(4) = (5) + (1, \text{Angus})$$

$$(6) = (4) + b[(2) - (3)] \text{ with } b = 1.14$$

$$(7) = (6) - (6, \text{Angus})$$

$$(8) = (7) - (7, \text{Angus}) - [(2) - (2, \text{Angus})]$$

Table 4. Breed of maternal grandsire solutions from MARC, mean breed and MARC EPD used to adjust for genetic trend to the year 2004 base and factors to adjust within-breed EPD to an Angus equivalent – MILK (lb)

| Breed | MGS | Number | | Raw | Mean EPD | | | | Breed Soln | Adjust to | | Factor to | | |
|-------------|-----|--------|-----------|---------------|---------------|---------------|---------|---------------|---------------------------------|-----------------------------------|-----------|-------------------------------|-------|-------|
| | | Gpr | Daughters | MARC Mean (1) | Breed WWT (2) | MARC MILK (3) | WWT (4) | MARC MILK (5) | at MARC MWWT Ang vs Ang (6) (7) | 2004 Base MWWT Ang vs Ang (8) (9) | MILK (10) | Adjust MILK EPD to Angus (11) | | |
| Hereford | 108 | 2890 | 714 | 482 | 37.0 | 14.0 | 22.7 | 7.1 | 482 | -20.2 | 498 | -25.3 | -17.1 | -18.3 |
| Angus | 104 | 2152 | 541 | 502 | 38.5 | 19.0 | 20.3 | 9.6 | 502 | 0.0 | 523 | 0.0 | 6.2 | 0.0 |
| Shorthorn | 22 | 251 | 69 | 527 | 13.0 | 2.4 | 8.4 | 5.8 | 525 | 23.2 | 524 | 0.9 | 4.6 | 15.0 |
| South Devon | 14 | 347 | 69 | 488 | 19.1 | 7.1 | -0.3 | 5.5 | 505 | 3.2 | 518 | -5.0 | -0.4 | 5.3 |
| Brahman | 40 | 880 | 216 | 522 | 14.1 | 6.1 | 4.5 | 2.7 | 533 | 31.1 | 542 | 19.3 | 19.8 | 26.5 |
| Simmental | 47 | 1129 | 239 | 522 | 34.1 | 5.4 | 22.6 | 8.0 | 527 | 24.8 | 530 | 7.3 | 4.5 | 11.9 |
| Limousin | 40 | 1112 | 240 | 489 | 36.3 | 18.3 | 19.8 | 15.6 | 493 | -8.7 | 506 | -17.3 | -10.9 | -16.4 |
| Charolais | 68 | 1029 | 235 | 510 | 20.0 | 6.2 | 7.4 | 3.7 | 512 | 9.7 | 522 | -1.3 | -5.2 | 1.3 |
| Maine-Anjou | 17 | 485 | 86 | 533 | 39.6 | 18.4 | 46.4 | 21.8 | 520 | 18.4 | 513 | -10.3 | -1.6 | -7.2 |
| Gelbvieh | 46 | 989 | 232 | 536 | 41.0 | 18.0 | 32.3 | 17.2 | 525 | 23.1 | 531 | 8.0 | 9.8 | 4.6 |
| Tarentaise | 6 | 341 | 78 | 513 | 4.0 | 1.0 | -5.9 | 4.6 | 521 | 18.6 | 522 | -0.8 | 6.8 | 18.6 |
| Salers | 25 | 351 | 89 | 534 | 15.5 | 8.7 | 5.5 | 11.9 | 525 | 22.9 | 527 | 3.9 | 6.1 | 11.2 |
| Red Angus | 21 | 330 | 89 | 505 | 29.0 | 15.0 | 27.6 | 13.7 | 503 | 1.1 | 505 | -17.7 | -5.9 | -8.1 |
| Braunvieh | 7 | 502 | 92 | 542 | 7.0 | 0.0 | 7.9 | -0.4 | 527 | 25.4 | 527 | 4.3 | 11.3 | 24.1 |
| Brangus | 19 | 99 | 43 | 527 | 22.7 | 9.9 | 24.7 | 4.7 | 509 | 6.6 | 513 | -9.6 | -5.5 | -2.6 |
| Beefmaster | 20 | 113 | 51 | 528 | 7.0 | 2.0 | 15.4 | -1.1 | 504 | 1.7 | 502 | -20.6 | -17.9 | -7.1 |

Calculations:

(6) = (7) + (1, Angus); (8) = (6) + $b_{WWT}[(2) - (4)] + b_{MLK} [(3) - (5)]$ with $b_{WWT} = 0.57$ and $b_{MLK} = 1.13$; (9) = (8) - (8, Angus); (10) = [(9) - Average (9)] - 0.5[(7, Table 2) - Average (7, Table 2)]; (11) = [(10) - (10, Angus)] - [(3) - (3, Angus)].

Table 5. Mean weighted^a accuracies for birth weight (BWT), weaning weight (WWT), yearling weight (YWT), maternal weaning weight (MWWT) and milk (MILK) for bulls used at MARC

| Breed | BWT | WWT | YWT | MWWT | MILK |
|-------------|------|------|------|------|------|
| Hereford | 0.56 | 0.53 | 0.48 | 0.51 | 0.50 |
| Angus | 0.72 | 0.75 | 0.69 | 0.72 | 0.64 |
| Shorthorn | 0.65 | 0.59 | 0.57 | 0.61 | 0.48 |
| South Devon | 0.37 | 0.39 | 0.37 | 0.41 | 0.42 |
| Brahman | 0.50 | 0.55 | 0.38 | 0.55 | 0.42 |
| Simmental | 0.94 | 0.94 | 0.93 | 0.94 | 0.93 |
| Limousin | 0.92 | 0.89 | 0.82 | 0.89 | 0.84 |
| Charolais | 0.71 | 0.65 | 0.56 | 0.65 | 0.56 |
| Maine-Anjou | 0.72 | 0.71 | 0.71 | 0.71 | 0.71 |
| Gelbvieh | 0.72 | 0.66 | 0.51 | 0.71 | 0.58 |
| Tarentaise | 0.95 | 0.95 | 0.95 | 0.95 | 0.95 |
| Salers | 0.83 | 0.83 | 0.77 | 0.82 | 0.83 |
| Red Angus | 0.87 | 0.85 | 0.84 | 0.84 | 0.81 |
| Braunvieh | 0.84 | 0.85 | 0.83 | 0.85 | 0.77 |
| Brangus | 0.80 | 0.78 | 0.65 | 0.81 | 0.64 |
| Beefmaster | 0.67 | 0.75 | 0.62 | 0.77 | 0.64 |

^aWeighted by number of progeny at MARC for BWT, WWT, and YWT and by number of grandprogeny for MWWT and MILK.

Table 6. REML estimates of variance components (lb^2) for birth weight (BWT), weaning weight (WWT), yearling weight (YWT), and maternal weaning weight (MWWT) from mixed model analyses

| Analysis | Direct | | | Maternal |
|---------------------------------|--------|---------|---------|----------|
| | BWT | WWT | YWT | MWWT |
| Direct ^a | | | | |
| Sires (650) within breed (17) | 11.42 | 152.19 | 630.84 | |
| Dams (4395) within breed (3) | 26.62 | 875.77 | 1232.73 | |
| Residual | 68.17 | 1534.80 | 4036.68 | |
| Maternal | | | | |
| MGS (619) within MGS breed (17) | | | | 210.79 |
| Daughters within MGS (3214) | | | | 966.90 |
| Residual | | | | 1378.92 |

^aNumbers for weaning weight.

Table 7. Pooled regression coefficients (lb/lb) for weights at birth (BWT), 205 days (WWT), and 365 days (YWT) of F₁ progeny on sire expected progeny difference and by sire breed, dam breed, and sex of calf

| | BWT | WWT | YWT |
|--------------------|-------------|--------------|--------------|
| Pooled | 1.02 ± 0.05 | 0.89 ± 0.05 | 1.14 ± 0.05 |
| Sire breed | | | |
| Hereford | 1.20 ± 0.08 | 0.87 ± 0.08 | 1.18 ± 0.07 |
| Angus | 0.85 ± 0.10 | 0.79 ± 0.10 | 1.15 ± 0.08 |
| Shorthorn | 0.52 ± 0.47 | 0.60 ± 0.46 | 1.23 ± 0.35 |
| South Devon | 0.91 ± 0.56 | -0.17 ± 0.36 | -0.04 ± 0.40 |
| Brahman | 1.82 ± 0.27 | 1.12 ± 0.27 | 0.68 ± 0.24 |
| Simmental | 1.02 ± 0.22 | 1.26 ± 0.17 | 1.31 ± 0.16 |
| Limousin | 0.69 ± 0.16 | 0.55 ± 0.16 | 1.16 ± 0.15 |
| Charolais | 1.02 ± 0.14 | 0.94 ± 0.13 | 0.91 ± 0.12 |
| Maine-Anjou | 1.01 ± 0.36 | 0.52 ± 0.48 | 0.23 ± 0.48 |
| Gelbvieh | 1.00 ± 0.16 | 1.25 ± 0.27 | 1.34 ± 0.22 |
| Tarentaise | 0.60 ± 0.85 | 0.75 ± 0.55 | 1.31 ± 0.59 |
| Salers | 1.19 ± 0.38 | 0.98 ± 0.45 | 0.77 ± 0.45 |
| Red Angus | 0.55 ± 0.19 | 0.60 ± 0.34 | 0.77 ± 0.30 |
| Braunvieh | 0.47 ± 0.37 | 0.87 ± 0.82 | 2.06 ± 0.54 |
| Brangus | 1.44 ± 0.34 | 0.81 ± 0.43 | 0.46 ± 0.40 |
| Beefmaster | 1.11 ± 0.52 | 1.43 ± 0.34 | 1.37 ± 0.38 |
| Dam breed | | | |
| Hereford | 0.91 ± 0.08 | 0.83 ± 0.08 | 1.01 ± 0.07 |
| Angus | 1.12 ± 0.06 | 0.93 ± 0.07 | 1.19 ± 0.06 |
| MARC III | 0.98 ± 0.08 | 0.88 ± 0.09 | 1.20 ± 0.08 |
| Sex of calf | | | |
| Heifers | 1.00 ± 0.06 | 0.99 ± 0.06 | 1.00 ± 0.06 |
| Steers | 1.04 ± 0.06 | 0.80 ± 0.06 | 1.28 ± 0.06 |

Table 8. Pooled regression coefficients (lb/lb) for progeny performance on maternal grandsire EPD for weaning weight (MWWT) and milk (MILK) and by breed of maternal grandsire, breed of maternal grandam, and sex of calf

| Type of regression | MWWT | MILK |
|------------------------------------|-------------|--------------|
| Pooled | 0.57 ± 0.04 | 1.13 ± 0.06 |
| Breed of maternal grandsire | | |
| Hereford | 0.50 ± 0.07 | 1.15 ± 0.11 |
| Angus | 0.53 ± 0.08 | 1.07 ± 0.12 |
| Shorthorn | 0.64 ± 0.39 | 0.03 ± 0.63 |
| South Devon | 0.31 ± 0.25 | -1.23 ± 0.85 |
| Brahman | 0.42 ± 0.21 | 0.51 ± 0.34 |
| Simmental | 0.91 ± 0.17 | 1.34 ± 0.40 |
| Limousin | 1.14 ± 0.12 | 1.87 ± 0.23 |
| Charolais | 0.39 ± 0.11 | 1.25 ± 0.20 |
| Maine-Anjou | 0.17 ± 0.34 | 0.48 ± 0.38 |
| Gelbvieh | 1.01 ± 0.22 | 1.48 ± 0.32 |
| Tarentaise | 0.27 ± 0.70 | 0.89 ± 0.85 |
| Salers | 0.89 ± 0.33 | 2.25 ± 0.37 |
| Red Angus | 0.94 ± 0.27 | 1.94 ± 0.30 |
| Braunvieh | 2.81 ± 1.15 | -0.17 ± 1.53 |
| Brangus | 0.06 ± 0.73 | 0.16 ± 0.68 |
| Beefmaster | 1.13 ± 0.50 | 3.92 ± 0.56 |
| Breed of maternal grandma | | |
| Hereford | 0.54 ± 0.06 | 1.55 ± 0.10 |
| Angus | 0.57 ± 0.05 | 1.05 ± 0.09 |
| MARC III | 0.59 ± 0.07 | 0.89 ± 0.10 |
| Sex of calf | | |
| Heifers | 0.57 ± 0.05 | 1.12 ± 0.08 |
| Steers | 0.57 ± 0.05 | 1.14 ± 0.08 |

Table 9. Variances (lb²) of adjusted breed differences to add to sum of within breed prediction error variances to obtain variance of differences of across breed EPD for bulls of two different breeds^a. Birth weight above the diagonal and yearling weight below the diagonal

| Breed | HE | AN | SH | SD | BR | SI | LI | CH | MA | GE | TA | SA | RA | BV | BS | BM |
|-------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| HE | 0.0 | 0.2 | 0.9 | 1.4 | 0.5 | 0.5 | 0.5 | 0.4 | 1.0 | 0.4 | 2.6 | 0.8 | 0.8 | 1.2 | 0.9 | 1.0 |
| AN | 14 | 0.0 | 0.9 | 1.4 | 0.5 | 0.5 | 0.5 | 0.4 | 1.1 | 0.5 | 2.6 | 0.8 | 0.8 | 1.2 | 0.9 | 1.0 |
| SH | 55 | 57 | 0.0 | 2.0 | 1.2 | 1.2 | 1.2 | 1.0 | 1.6 | 1.0 | 3.2 | 1.2 | 1.5 | 1.8 | 1.7 | 1.8 |
| SD | 83 | 84 | 125 | 0.0 | 1.7 | 1.3 | 1.4 | 1.3 | 2.1 | 1.6 | 3.7 | 1.9 | 1.8 | 2.3 | 2.2 | 2.2 |
| BR | 36 | 37 | 81 | 110 | 0.0 | 0.9 | 0.9 | 0.8 | 1.3 | 0.8 | 2.6 | 1.1 | 1.1 | 1.5 | 1.3 | 1.4 |
| SI | 28 | 29 | 72 | 80 | 56 | 0.0 | 0.5 | 0.5 | 1.3 | 0.6 | 2.8 | 1.1 | 0.8 | 1.4 | 1.3 | 1.3 |
| LI | 31 | 32 | 75 | 83 | 58 | 30 | 0.0 | 0.5 | 1.3 | 0.7 | 2.9 | 1.1 | 0.8 | 1.5 | 1.3 | 1.3 |
| CJ | 24 | 25 | 63 | 81 | 52 | 28 | 31 | 0.0 | 1.2 | 0.5 | 2.7 | 0.9 | 0.8 | 1.3 | 1.2 | 1.2 |
| MA | 62 | 64 | 100 | 128 | 86 | 75 | 78 | 71 | 0.0 | 1.0 | 3.2 | 1.5 | 1.6 | 1.1 | 1.9 | 1.9 |
| GE | 28 | 30 | 66 | 95 | 54 | 37 | 39 | 34 | 62 | 0.0 | 2.8 | 0.9 | 0.8 | 1.2 | 1.2 | 1.3 |
| TA | 151 | 154 | 190 | 220 | 158 | 167 | 170 | 163 | 191 | 164 | 0.0 | 3.1 | 3.2 | 3.4 | 3.4 | 3.4 |
| SA | 48 | 51 | 73 | 118 | 74 | 66 | 68 | 57 | 93 | 60 | 184 | 0.0 | 1.4 | 1.7 | 1.6 | 1.7 |
| RA | 46 | 46 | 91 | 111 | 75 | 49 | 51 | 48 | 95 | 52 | 188 | 84 | 0.0 | 1.7 | 1.5 | 1.5 |
| BV | 69 | 72 | 108 | 135 | 93 | 83 | 85 | 79 | 67 | 69 | 198 | 101 | 102 | 0.0 | 2.0 | 2.1 |
| BS | 64 | 65 | 116 | 141 | 96 | 85 | 87 | 82 | 122 | 85 | 212 | 109 | 99 | 129 | 0.0 | 1.0 |
| BM | 65 | 65 | 116 | 141 | 96 | 85 | 87 | 82 | 122 | 85 | 212 | 109 | 100 | 129 | 76 | 0.0 |

^aFor example, a Hereford bull has within breed PEV of 300 for YWT and that for a Shorthorn bull is 200. Then the PEV for the difference in EPDs for the two bulls is $55 + 300 + 200 = 555$ with $SEP = \text{SQRT}(555) = 23.6$.

Table 10. Variances (lb²) of adjusted breed differences to add to sum of within breed prediction error variances to obtain variance of difference of across breed EPDs for bulls of two different breeds. Weaning weight direct above the diagonal and MILK below the diagonal

| Breed | HE | AN | SH | SD | BR | SI | LI | CH | MA | GE | TA | SA | RA | BV | BS | BM |
|-------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|
| HE | 0 | 4 | 20 | 28 | 11 | 9 | 10 | 8 | 22 | 9 | 42 | 18 | 17 | 24 | 20 | 20 |
| AN | 13 | 0 | 21 | 28 | 11 | 10 | 10 | 8 | 23 | 9 | 43 | 18 | 17 | 25 | 20 | 20 |
| SH | 50 | 52 | 0 | 44 | 28 | 26 | 27 | 23 | 37 | 24 | 57 | 27 | 34 | 39 | 39 | 39 |
| SD | 59 | 60 | 99 | 0 | 36 | 27 | 28 | 27 | 45 | 32 | 66 | 42 | 39 | 47 | 46 | 46 |
| BR | 26 | 27 | 67 | 77 | 0 | 18 | 18 | 16 | 29 | 16 | 43 | 25 | 26 | 31 | 29 | 29 |
| SI | 25 | 26 | 65 | 61 | 43 | 0 | 10 | 9 | 27 | 12 | 48 | 24 | 18 | 29 | 27 | 27 |
| LI | 27 | 28 | 67 | 63 | 44 | 30 | 0 | 10 | 28 | 13 | 48 | 25 | 18 | 29 | 28 | 27 |
| CJ | 21 | 22 | 58 | 60 | 38 | 27 | 29 | 0 | 26 | 11 | 46 | 21 | 18 | 27 | 26 | 26 |
| MA | 56 | 58 | 93 | 103 | 72 | 69 | 71 | 65 | 0 | 22 | 58 | 35 | 35 | 24 | 41 | 41 |
| GE | 23 | 24 | 59 | 69 | 40 | 34 | 35 | 29 | 59 | 0 | 46 | 21 | 19 | 23 | 27 | 27 |
| TA | 126 | 128 | 164 | 175 | 130 | 141 | 143 | 136 | 167 | 137 | 0 | 55 | 56 | 59 | 60 | 60 |
| SA | 42 | 44 | 70 | 91 | 59 | 57 | 59 | 50 | 85 | 51 | 156 | 0 | 31 | 37 | 36 | 37 |
| RA | 45 | 45 | 85 | 88 | 62 | 50 | 51 | 48 | 89 | 50 | 161 | 77 | 0 | 37 | 34 | 34 |
| BV | 82 | 84 | 120 | 130 | 98 | 96 | 97 | 91 | 97 | 85 | 194 | 112 | 116 | 0 | 42 | 42 |
| BS | 75 | 76 | 119 | 127 | 94 | 93 | 94 | 89 | 125 | 91 | 195 | 111 | 111 | 151 | 0 | 21 |
| BM | 67 | 68 | 111 | 119 | 86 | 85 | 86 | 81 | 117 | 83 | 187 | 103 | 103 | 143 | 103 | 0 |