


Economic Benefits of using Genetic Selection to Reduce the Prevalence of Bovine Respiratory Disease Complex in Beef Feedlot Cattle

Bovine Respiratory Disease Complex Coordinated Agriculture Project




<http://www.brdcomplex.org/>

H.L. Neibergs¹, J.S. Neibergs¹, A.J. Wojtowicz¹, J.F. Taylor², C.M. Seabury³, J. E. Womack³

¹Washington State University, Pullman, WA; ²University of Missouri, Columbia, MO; ³Texas A&M University, College Station, TX

BOVINE RESPIRATORY DISEASE COMPLEX




- Most common infectious disease of cattle worldwide
- Leading cause of death in dairy and beef cattle (NASS 2006)
- Despite use of vaccines and antimicrobials, morbidity and mortality rates have remained virtually unchanged in cattle over the past several decades (Miles 2009; Gagea et al. 2006)

WASHINGTON STATE UNIVERSITY

BOVINE RESPIRATORY DISEASE COMPLEX


- The average annual prevalence of BRDC was 16.2% over a 15 year period (USDA 2011)
- ~97% of feedlots with ≥ 1000 head of cattle have BRDC (USDA 2011)
- 60% of all harvested feedlot cattle have lung lesions from BRDC (Schneider et al 2009)



WASHINGTON STATE UNIVERSITY

ECONOMICS OF BRDC

- BRDC remains the most economically important disease of beef cattle, responsible for losses of over \$800 million annually (Chirase & Greene 2001; Snowden et al. 2006; USDA 2001; Gagea et al. 2006)
- Approximately 350,000 feedlot cattle die annually in the US due to BRDC (USDA 2011)



WASHINGTON STATE UNIVERSITY

PROBLEM

Same level of morbidity and mortality from BRDC over the past 20 years despite utilizing:


- Best management practices
 - Preventative vaccines
 - Improved treatments

We need new approaches to reduce the incidence of BRDC in addition to our current approaches that are economically feasible!

WASHINGTON STATE UNIVERSITY

GENETICS


- There is increasing evidence that susceptibility to BRDC is at least partly under genetic control
- Differences in BRDC susceptibility has been found between cattle breeds and sire lines (Muggli-Cockett et al. 1992; Snowden et al. 2006; Maltecca et al. 2006)



WASHINGTON STATE UNIVERSITY

GENETICS

- Heritability estimates for susceptibility to BRDC have ranged from 0% to 26% in beef and dairy cattle (Snowder et al. 2005, 2006; Muggli-Cockett et al. 1992; Heringstad et al. 2008, Seabury et al. 2014, Neibergs et al. 2014)
- Studies in crossbred beef cattle identified loci associated with susceptibility to BRDC and cattle persistently infected with bovine viral diarrhea virus (Zanella et al. 2011)



Animals & Phenotypes



FEEDLOT STEERS



- Samples collected on 995 beef steers: 908 Angus, 18 Charolais, 25 Hereford, 44 Red Angus
- Cases and controls were housed in the same pens
- 497 were affected with BRDC, 498 were unaffected
- Animal health status defined by fever, cough, nasal discharge, eye discharge, and ear position or head tilt



SCORING RESPIRATORY DISEASE

Calf Health Scoring Criteria				
	0	1	2	3
Rectal temperature	100-100.5	101-101.5	102-102.5	>103
Cough	None	Induced single cough	Induced repeated coughs or occasional spontaneous coughs	Repeated spontaneous coughs
Nasal discharge	Normal serous discharge	Small amount of unilateral cloudy discharge	Bilateral, cloudy or excessive mucous discharge	Copious bilateral mucopurulent discharge
Eye scores	Normal	Small amount of ocular discharge	Moderate amount of ocular discharge	Heavy ocular discharge
Ear scores	Normal	Ear flick or head shake	Slight unilateral droop	Head tilt or bilateral droop

McGuirk SM. 2008. Disease management of dairy calves and heifers. Vet Clin NA: Food Anim Pract 24:139-153.

DIAGNOSTICS AND GENOTYPING

- Diagnostics for *Mycoplasma*, *P. Multocida*, *M. Haemolytica*, *H. Somni*, bovine respiratory syncytial virus, bovine viral diarrhea virus, IBR (bovine herpes virus 1) completed
- DNA extracted, and genotyped for 778,000 SNPs



FEEDLOT STEERS

- Data on lung and liver scores, yield grade, quality grade, and condemned carcasses were collected at harvest
- Treatment costs, initial weights, hot carcass weight, days on feed, collected for both cases and controls
- Pens were harvested as a group





RESULTS

HEALTH SCORES

- Mean clinical score for cases was 8.04 ± 1.23
- Mean clinical score for controls was 2.06 ± 0.037
- 67.7% of cases had lung lesions
- 67.2% of controls had lung lesions
- Mortality for cases was 5.3% and 0.4% for controls



DIAGNOSTIC SWAB RESULTS



Washington Animal Disease Diagnostic Lab

Pathogen	Feedlot Positive Cases	Feedlot Positive Controls	Odds Ratio	Odds Ratio 95% Confidence Interval	Odds Ratio P value
<i>Arcanobacterium pyogenes</i>	3.4%	0.8%	4.35	1.41-17.89	0.0066
<i>Histophilus somni</i>	26.2%	12.7%	2.41	1.72-3.41	<0.0001
<i>Mannheimia haemolytica</i>	38.2%	22.5%	2.11	1.59-2.82	<0.0001
<i>Pasteurella multocida</i>	36.4%	36.1%	1.01	0.79-1.31	1.0 (*NS)
<i>Mycoplasma</i> spp.	84.4%	77.9%	1.54	1.1-2.15	<0.0097
Bovine corona virus	17.4%	9.6%	2.08	1.4-3.11	0.0001
Bovine respiratory syncytial virus	2.4%	0.8%	2.53	0.82-9.24	0.0881 (*NS)
Bovine viral diarrhea virus	4.0%	1.6%	2.79	1.16-7.41	0.0125
Bovine herpes virus	3.2%	1.6%	2.09	0.83-5.7	0.0995 (*NS)

*NS- not significant

HERITABILITY ESTIMATES







- Heritability estimates were obtained by GenABEL/GRAMMAR from relationship matrixes obtained from the genotypes of the BovineHD assay
- Heritability estimates were **17.7%** for BRDC as a case-control and **29.2%** for BRDC as a categorical trait

RATE OF GENETIC CHANGE

Factors	Case-Control	Health Score
BRDC prevalence (USDA 2011)	16.2%	16.2%
Estimated heritability	17.7%	29.2%
Accuracy of selection	0.42	0.54
Selection intensity	1.16	1.16
Additive genetic variance	0.15	0.19
Generation interval (years)	6	6
Rate of Genetic Change	1.26%	2.08%



ECONOMICS



QUALITY GRADE				
	Cases	Controls	χ^2	Significance
Prime	16	12	1.3	NS
Choice	240	286	8.1	0.005
Select	181	177	0	NS
No rolls	10	8	0.5	NS
Condemned	2	0	NA	NS
Deads	26	2	285.3	<0.0001
Railers	17	3	64.6	<0.0001
Total	492	488		



MARBLING SCORES				
	Quality	Case	Control	Significance
P=0.009	Prime+	0	0	NA
	Prime Avg	0	1	NS
	Prime -	7	4	NS
	Choice+	17	19	NS
	Choice Avg	44	74	0.001
	Choice-	179	200	NS
	Select+	134	127	NS
	Select -	49	44	NS
	Standard+	7	4	NS
	Standard-	12	10	NS
		449	483	

FEEDLOT COST OF BRDC
<ul style="list-style-type: none"> Average loss in carcass value (and death loss) was \$162.78, the cost of a single Draxxin treatment was \$41.32, average feeder purchase price lost (of those that died) across all cases was \$49.87 \$253.97 was the total cost per BRDC affected animal If CAB premiums were included, the cost would be higher
WASHINGTON STATE UNIVERSITY

POTENTIAL GAINS
<ul style="list-style-type: none"> In 2013, 9,131,500 heifers and 16,003,400 steers were harvested from US feedlots with >1000 head of cattle 4,071,854 feedlot cattle were estimated to be affected with BRDC with a 16.2% prevalence rate When feeder purchase costs are included, the cost of BRDC was \$253.97 per affected animal or \$1,034,128,760 in 2013
<i>This is a conservative estimate</i>
WASHINGTON STATE UNIVERSITY

POTENTIAL GAINS
<p>With the estimated rate of genetic gain for selection for BRDC at 1.26% (case-control) and 2.08%, between \$13-\$21.5 million could have been saved in the feedlot sector through selection in 2013</p>

ACKNOWLEDGEMENT'S
<p>Bovine Respiratory Disease Complex Coordinated Agricultural Project</p>
<p>http://www.brdcomplex.org/</p>
<p>Funded by: National Research Initiative competitive Grant no. 2011-68004-30367 from the USDA National Institute of Food and Agriculture</p>

